



## Iliotibial Band Syndrome Rehabilitation Exercises



Iliotibial band stretch (standing)



Iliotibial band stretch (side-leaning)



Standing calf stretch



Hamstring stretch on wall



Quadriceps stretch



Wall squat with a ball


# Iliotibial Band Syndrome Rehabilitation Exercises



Side-lying leg lift




Knee stabilization: A



Knee stabilization: B



Knee stabilization: C



Knee stabilization: D



Iliotibial band stretch (side-bending)



Clam exercise

## Iliotibial Band Syndrome Rehabilitation Exercises

You may do all of these exercises right away.

- **Iliotibial band stretch: Standing:** Cross one leg in front of the other leg and bend down and touch your toes. You can move your hands across the floor toward the front leg and you will feel more stretch on the outside of your thigh on the other side. Hold this position for 15 to 30 seconds. Return to the starting position. Repeat 3 times. Reverse the positions of your legs and repeat.
- **Iliotibial band stretch: Side-leaning:** Stand sideways near a wall. Place one hand on the wall for support. Cross the leg farthest from the wall over the other leg, keeping the foot closest to the wall flat on the floor. Lean your hips into the wall. Hold the stretch for 15 seconds, repeat 3 times, and then switch legs and repeat the exercise another 3 times.
- **Standing calf stretch:** Facing a wall, put your hands against the wall at about eye level. Keep one leg back with the heel on the floor, and the other leg forward. Turn your back foot slightly inward (as if you were pigeon-toed) as you slowly lean into the wall until you feel a stretch in the back of your calf. Hold for 15 to 30 seconds. Repeat 3 times and then switch the position of your legs and repeat the exercise 3 times. Do this exercise several times each day.
- **Hamstring stretch on wall:** Lie on your back with your buttocks close to a doorway, and extend your legs straight out in front of you along the floor. Raise one leg and rest it against the wall next to the door frame. Your other leg should extend through the doorway. You should feel a stretch in the back of your thigh. Hold this position for 15 to 30 seconds. Repeat 3 times and then switch legs and do the exercise again.
- **Quadriceps stretch:** Stand an arm's length away from the wall with your injured leg farthest from the wall. Facing straight ahead, brace yourself by keeping one hand against the wall. With your other hand, grasp the ankle of your injured leg and pull your heel toward your buttocks. Don't arch or twist your back. Keep your knees together. Hold this stretch for 15 to 30 seconds.
- **Wall squat with a ball:** Stand with your back, shoulders, and head against a wall and look straight ahead. Keep your shoulders relaxed and your feet 2 feet away from the wall and a shoulder's width apart. Place a soccer or basketball-sized ball behind your back. Keeping your back upright, slowly squat down to a 45-degree angle. Your thighs will not yet be parallel to the floor. Hold this position for 10 seconds and then slowly slide back up the wall. Repeat 10 times. Build up to 3 sets of 10.
- **Side-lying leg lift:** Lying on your uninjured side, tighten the front thigh muscles on your top leg and lift that leg 8 to 10 inches away from the other leg. Keep the leg straight and lower slowly. Do 3 sets of 10.
- **Knee stabilization:** Wrap a piece of elastic tubing around the ankle of the uninjured leg. Tie a knot in the other end of the tubing and close it in a door.
  1. Stand facing the door on the leg without tubing and bend your knee slightly, keeping your thigh muscles tight. While maintaining this position, move the leg with the tubing straight back behind you. Do 3 sets of 10.
  2. Turn 90 degrees so the leg without tubing is closest to the door. Move the leg with tubing away from your body. Do 3 sets of 10.
  3. Turn 90 degrees again so your back is to the door. Move the leg with tubing straight out in front of you. Do 3 sets of 10.

4. Turn your body 90 degrees again so the leg with tubing is closest to the door. Move the leg with tubing across your body. Do 3 sets of 10.

Hold onto a chair if you need help balancing. This exercise can be made even more challenging by standing on a pillow while you move the leg with tubing.

- **Iliotibial band stretch: Side-bending:** Cross one leg in front of the other leg and lean in the opposite direction from the front leg. Reach the arm on the side of the back leg over your head while you do this. Hold this position for 15 to 30 seconds. Return to the starting position. Repeat 3 times and then switch legs and repeat the exercise.
- **Clam exercise:** Lie on your uninjured side with your hips and knees bent and feet together. Slowly raise your top leg toward the ceiling while keeping your heels touching each other. Hold for 2 seconds and lower slowly. Do 3 sets of 10 repetitions.

Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for RelayHealth.

Published by RelayHealth.

© 2009 RelayHealth and/or its affiliates. All Rights Reserved.